


HELP YOUR PET HEAL!

for the next 7-10 days

Watch a
video here



NO LICKING

Licking or chewing at the incision area can cause infection.




INSPECT

What you see today is normal. Check twice a day for redness, swelling or discharge.

KEEP IT DRY



Water affects the surgical glue and can cause the incision to open.



SEPERATE ANIMALS

Keep females away from unneutered males.



MALES



Incisions are left open & it is normal to see swelling and red discharge for a few days.

RESTRICT ACTIVITY



Inside rest gives the tissue time to heal & helps the incision to stay closed.
NO RUNNING, JUMPING OR PLAYING!



FEED A SIMPLE DIET

No junk food, table scraps, milk or extra treats. The anesthesia can upset their tummy.

NO HUMAN MEDICATION

It can be fatal! Your pet has already been given a long-acting pain medication.



If these instructions are followed, in full, we will make every reasonable effort to treat at our clinic, at minimal cost, any postoperative complications resulting directly from surgery. Your regular veterinarian must address illnesses or injuries that are not a direct result of surgery. We are available after hours but cannot be available 24/7. Your pet is your responsibility, please do not withhold emergency treatment, if needed, because you cannot reach us.

Critter Fixers-
Emergency Line

**615
389
4399**

Contact us immediately if you notice: Pale gums; depression; problems walking; loss of appetite or decreased water intake; vomiting; diarrhea; heavy discharge or bleeding from the incision; difficulty with urination or bowel movements; difficulty breathing.